

09875935.060801

A. Most strongly disease-associated patterns								
Marker	01	02	03	04	05	06	07	08
Pattern01	4	1	6	6	3	*	*	*
Pattern02	4	1	6	6	3	6	*	*
Pattern03	4	1	6	6	*	6	*	*
Pattern04	4	1	6	*	3	6	*	*
Pattern05	4	1	*	6	3	*	*	*
Pattern06	4	1	*	6	3	6	*	*
Pattern07	4	*	6	6	3	*	*	*
Pattern08	4	*	6	6	3	6	*	*
Pattern09	4	*	6	*	3	6	*	*
Pattern10	*	1	6	6	*	*	*	*
Pattern11	*	1	6	6	3	*	*	*

Frequency	9	11	11	11	10	6	0	0

FIG. 1A

B. Marker frequency histogram




FIG 1B

C. Marker frequency histogram




FIG. 1C

D. Prediction results (A = 10%)




FIG. 1D

A. Influence of A




FIG. 2A

B. Influence of A (larger dataset)




FIG. 2 B

C. Influence of corrupted data ($A = 10\%$)




FIG. 2C

09875935.060801
T08090" 5E657860

D. Influence of missing data (A = 10%)




FIG. 2D

09052652360

E. Comparison of prediction methods
(A = 10%, 0/1% corrupted, 0/20% missing)




FIG. 2E

F. Influence of pattern parameters
(A = 10%, 1% corrupted, 20% missing)




FIG. 2F

09875935.060801




FIG. 3A

B. Marker-wise frequencies for some p values




FIG. 3B

T08090" SE65/860

C. Marker-wise frequencies for some p values




FIG. 3C

D. Influence of permutation tests ($A = 5\%$)




FIG. 3D

Influence of A (SNP dataset, 12.5% missing data)




FIG. 4

A. Marker frequencies in HLA data




FIG. 5A

09875935-050801
T08090"SE65Z860

B. P values and frequency ratios in HLA data




FIG. 5B

A. Marker frequency histogram (2 genes)




FIG. 6A

005595.0001




FIG. 6B

T08090" SE654860

C. Localization residuals (2 genes)




FIG. 6C

09875935.060804
T08090" SE654860

D. Prediction accuracy (2 genes)




FIG. 6D

T00090" 5654860




FIG. 7A




FIG. 7B